

Transit ReLeaf®

ReLeaf® is a patented product formulation designed to eradicate biotic and abiotic stresses in high yielding and sensitive crops in commercial agriculture.

Elements	Typical Analysis (W/V)
Biologically Active Patented Formulation	32.8%
Organically Derived Amino Acids x16	48.2%
Other Organic Acids	8.2%

ReLeaf® can be applied to most horticultural crops and broad-acre crops with agronomic advice.

ReLeaf® can be used with fertilizers, biological materials, hormones and non-selective herbicides.

COMPATIBILITY: Always perform a compatibility test before mixing anything with ReLeaf®.

CAUTION: DO NOT MIX ReLeaf® with product containing Copper.

FOLIAR APPLICATION RATES

Apply ReLeaf® at 1 - 1.5ml/1L of water mixed with Weedicide/Herbicide on weeds for weed control.

Apply ReLeaf® at 1 - 1.5ml/1L of water and spray on to the foliage with recommended rate of fungicide/insecticide; repeat as needed.

Example of mixing:

Method 1: If your water rate for weedicide application is 300L/ha, you can mix ReLeaf® 300ml - 450ml per ha.

Method 2: If your weedicide tank volume is 2000L, you can add ReLeaf® 2000ml (2L) - 3000ml (3L) to the filled weedicide tank.

SOIL APPLICATION RATES

ReLeaf® can be applied with almost all farm chemicals and fertilizers.

ALL FIELD CROPS (grain crops, pastures, flowers, etc): Apply 500ml - 1L per hectare of ReLeaf®. Repeat the application every 6 - 8 weeks.

ALL VEGETABLE CROPS: Apply 500ml - 1L per hectare of ReLeaf® before flowering and repeat every 12 - 16 days.

POTATOES & CARROTS: Apply 500ml - 1L per hectare of ReLeaf® at any stage in the season.

GREENHOUSE CROPS/HYDROPONICS: Apply 1 - 1.5ml/L of water ReLeaf® at any stage in the season. ReLeaf® can be applied to Hydroponic tanks at a rate of 500ml/1000L in tank A and same rate in tank B. Best to check pH of water before applying ReLeaf® and apply product accordingly. LIMITED WARRANTY.

APPLES, POME & STONE FRUIT: Apply 500ml - 1L per hectare of ReLeaf® at any stage in the season.

FRUIT TREES & AVOCADOS: Apply 500ml - 1L per hectare of ReLeaf® at any stage in the season.

KIWIFRUIT, TABLE GRAPES, WINE GRAPES: Apply 500ml - 1L per hectare of ReLeaf® at any stage in the season.

ReLeaf® is best to apply during the active growth stage. ReLeaf® may be used at any time throughout the season to increase the efficiency. However, the product absorption efficiency may differ according to the crop, weather factors and other agronomic practices. Manufacturer, distributor or reseller makes no warranty, whether expressed or implied concerning the use of this product other than for the purposes indicated on the label. Neither the manufacturer nor reseller shall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label. For best results and more specific recommendations, contact your nearest qualified agronomist or distributor.

SHAKE WELL BEFORE USE

BATCH NO:

IN A TRANSPORT EMERGENCY
DIAL 000 POLICE OR
FIRE BRIGADE

FOR SPECIALIST ADVICE IN AN
EMERGENCY DIAL 1 800 033 111
24 HOURS AUSTRALIA WIDE



WARNING

KEEP OUT OF REACH OF CHILDREN

Product Code DCTRL007 **Net 1000L**

DUAL CHELATE FERTILIZER PTY LTD IS AN
ISO 9001:2015
CERTIFIED COMPANY

Manufactured by:
Dual Chelate Fertilizer
162 New Guinea Road,
Robinvale Victoria 3549

info@dualchelate.com
dualchelate.com.au

