

Assess the Effectiveness of Zinc Fertilizers and Organic Activators on Germination and Early Seedling Growth in Lentil

Dual Chelate Fertilizer Pty Ltd, 162 New Guinea Road, Robinvale, Victoria 3540, Australia
Corresponding author: Tel: +61 3 5026 4052, Email: info@dualchelate.com

Abstract

Lentils play a crucial role in Australian grain production. Different internal and external factors affect seed germination and seedling growth. Nutrients play a crucial role in seed germination and seedling growth and development. This trial was conducted to analyze the impact of some Zinc fertilizers and organic activators on improving lentil seed germination and seedling growth in early stages. It was found that lentil seeds treated with Transit Zn and Momentum ZBM Trio significantly improved the shoot height. In addition, Transit Zn significantly increased the lentil root length. However, there was no significant difference between treatments in seed germination. Therefore, it can be recommended that the application of Transit Zn as well as Momentum ZBM Trio as seed treatments is highly beneficial in terms of improving seed germination and seedling growth.

Keywords: Lentil, Germination, Seedling growth, Root growth, Zinc Fertilizers, Organic Activators

1. Introduction

The grain industry is the Australian second largest agricultural industry which contributes 27% of the gross value of production (Our industry, 2022). Lentil is an important pulse crop in the world. Australia is one of the major lentil producers in the world and contributes to around 10% of the world's lentil production. Yield and the quality of the lentil yield are highly dependent on macro and micronutrient management (Sadras et al., 2021).

Seed germination is a fundamental process which affected by different external or environmental factors such as temperature, pH, and soil moisture and external factors as well as internal factors such as phytohormones, nutrients, etc. In addition, seed germination is regulated by different genes (Koger, Reddy, & Poston, 2004, Xue et al., 2021). Macro and micronutrients play a crucial role in seed germination, plant growth, and development. Zinc and phosphorus

are the key nutrients for better crop growth and yield (Arshad *et al.* 2016). Phosphorus and Zinc are readily absorbed leaves allowing rapid uptake and optimizing Zn and P levels at crucial early growth stages before soil-applied P is available. Moreover, Adequate delivery of Zinc & Phosphorus increases the production of energy molecules, and plant growth hormones and facilitates photosynthesis and nutrient transport. Zinc seed treatments have a great impact on improving seed germination, seedling growth, and development (Montanha et al., 2020). Prom-u-thai, Rerkasem, Yazici, & Cakmak found that seed priming with Zinc has a great influence on improving seed germination, root development, and dry weight in rice.

Zinc is a critical element for better and more vigorous wheat growth. It is both an activator and component of many enzymes and also influences auxin development (plant growth hormone) which promotes strong crop growth. Numerous studies have demonstrated that Zinc is responsible for a higher and a quality wheat yield

(Arshad *et al.* 2016). It is well documented that the Zinc content in seeds is highly affected by seed germination and seedling vigor in different broadacre crops such as wheat and barley (Imran, Mahmood, Neumann, & Bolt, 2021). Also, wheat seeds with high Zinc content can develop more shoots and roots at the early seedling stages and facilitates the uptake of more Zn in Zn-deficient soil (Graham & Rengel, 1993). Numerous studies have proven the impact of Zn seed treatment on seed germination, seedling growth, and yield. Harris, Rashid, Miraj, Arif, & Shah revealed that the seed treatment with 1% Zinc Sulphate improved the plant growth and yield in maize grown in Zn deficiency soil.

Dual Chelate Fertilizer Pty Ltd has developed a premium quality Zn fertilizer product called Momentum ZnP, Momentum ZBM Trio, Transit Zn as well as some other organic activators such as Transit Re-Leaf, Amino Boost Transit Max, and CPPA. Amino Boost Transit Max assists in increasing root growth, improving the translocation of nutrients, and enhancing the establishment of young plants. Amino Boost Transit Max® (ABTM) contains 10% Amino Acids, 6% Kelp, 4% Fulvic Acids, 1.5% organic activators, and 1.4% Nitrogen (Amino Acid derived Nitrogen). Momentum ZnP is a plant-available liquid Zinc & Phosphorus fertilizer designed to provide plants with optimal nutrients to promote early crop growth and establishment. Momentum ZnP consists of 18% Phosphorous, 14% Zinc, 2% Potassium, and patented organic activators. CPPA (Complex Polymeric Polyhydroxy Acid) is a group of organic acids that enhance various plant physiological functions such as nutrient absorption, shoot, and root growth. In this study, lentil seeds were treated with a 10% solution of each fertilizer product and organic activators to study the effectiveness of each product on seed germination and seedling growth.

- Measure the Shoot length and root length in each treatment
- Assess the impact of different fertilizer products on seedling growth and development.
- Visually compare the treatments to see any difference in appearance.
- Compare the benefits of treatments for lentil seed germination and seedling growth.

3. Materials and Methods

This trial was conducted in the greenhouse located at Robinvale. There were 8 treatments and 4 replicates in this trial. Hurricane lentil variety was selected for this trial and seeds were treated with 10% solution of each fertilizer product. Seeds were then placed in Petri dishes and each petri dish had 10 seeds. Petri dishes were misted, covered with lids, and then placed in a dark area. Germination count was recorded daily starting from day 3. Seedling measurements were taken after 12 days of treatment application.




Figure 1: Lentil seeds; variety Hurricane

2. Objectives

The specific objectives of this trial were to:

- Study the impact of treatments for seed germination.

Table 1: Application rates and product analysis of Zinc fertilizers and other organic activators.

Treatment	Rate	Product Analysis
Control	N/A	N/A
Amino Boost Transit Max	10%	10% Amino Acids, 6% Kelp, 4% Fulvic Acids, 1.5% Organic activators, 1.4% N
Momentum ZnP	10%	18.10% P, 2% K, 14% Zn
Momentum ZnP with Amino Acid and CPPA	10%	18.10% P, 2% K, 14% Zn + Amino Acid + CPPA
Momentum ZBM Trio	10%	5.30% B, 5.01% Zn, 0.24% Mo, 0.15% Mg + Amino acids
Transit Re-Leaf	10%	Fulvic acids, Amino acids, CPPA
Transit Zn	10%	3.67% N, 10.40% Zn, CPPA + Amino acids
CPPA	10%	Patented Organic

4. Observations

Germination Count

From day 3 onwards, the germination count was recorded daily and comparative photos were taken to visualize the impact of Zn products and organic activators on seed germination.

Shoot Height

After 12 days of seed sowing, seedlings were uprooted carefully, and measured the shoot height by using a ruler.

Root Length

After 12 days of seed sowing, seedlings were uprooted carefully, and measured the root length by using a ruler.

Comparative Photos

Comparative photos were taken to visually compare the seedling growth and root development.

5. Results




Figure 2: Visual comparison of seed germination in different treatments in day 4




Figure 3: Effects of different Zn fertilizer products and organic activators on Lentil seed germination in day 3.





Figure 4: Effects of different Zn fertilizer products and organic activators on Lentil shoot length. Asterisks represent statistical significance (* p,0.1, ** p,0.01).






Figure 5: Effects of different Zn fertilizer products and organic activators on Lentil root length. Asterisks represent statistical significance (***) p,0.001).



Figure 6: Visual comparison of seedling shoot and root growth in different treatments in day 12.

6. Discussion

Germination data were collected from day 3 of the seed sowing and all treatments had around 100% seed germination. Therefore, no significant difference was observed in germination counts (Figure 3). After 12 days of seed sowing, seedlings were uprooted, and measured the shoot height and root length. Transit Zn and Momentum ZBM Trio had the significantly highest shoot length compared to all other treatments. Momentum ZnP, Momentum ZnP with Amino acids & CPPA, and Transit Re-Leaf had the highest shoot height compared to the control seedlings, however, that was not a significant difference (Figure 4). After 12 days, Transit Zn showed the significantly highest root length compared to the control. Furthermore, Momentum ZnP, Momentum ZnP with amino acids and CPP, Momentum ZBM Trio, Transit Re-Leaf, and CPPA had the highest root length compared to the control seedlings (Figure 5). These results can be explained by the impact of Zinc, Phosphorous, Amino acids, and CPPA on seed germination as well as plant growth and development. Zinc is a crucial micronutrient for better and vigorous wheat growth. It is both an activator and component of many enzymes and also influences auxin development (plant growth hormone) which promotes

strong crop growth (Begum et al., 2016). Numerous studies have demonstrated that Zinc is responsible for a higher and a quality wheat yield (Arshad et al. 2016). Similarly, Auxin promotes stem elongation and guides shoot tips toward light sources which is a movement known as phototropism. Auxin also plays a role in maintaining apical dominance which explains the significant increment in plant growth parameters between the treatments and the control.

Furthermore, the application of Phosphorous is greatly influenced by yield maximizing in wheat (Grant and Baile, 1989). Phosphorous is the key nutrient for better root and shoot growth, especially in the early stages (Boring et al., 2018). Moreover, Phosphorus is incorporated into many organic compounds such as DNA, proteins, lipids, and enzymes. These organic compounds assist in energy transfer, nutrient uptake, and transport. A slow-release form of phosphorus allows for better nutrient utilization and absorption during the season (Talboys et al., 2015). Therefore, the increment of plant growth parameters in treated lentil plants should be due to the role of Phosphorous and Zinc. In addition, Zinc and Phosphorous have a great impact on plant root growth and several studies have demonstrated the importance of these nutrients on root growth in different plants such as *Zea mays L.* and rice (Hajabbasi and Schumacher, 1994, Phuphong et al., 2020). Therefore, this difference should be due to the influence of Zinc on seed germination and photosynthesis (OHKI, 1976).

Amino acids play a crucial role in plant growth and development. There are numerous studies have been conducted to assess the importance of amino acids on plant growth and development. Amino acids can, directly and indirectly, influence plant growth and development by affecting plant physiological activities. It was found that the foliar application of amino acid is beneficial for vegetative and reproductive growth as well as the yield quality of grapes (Khan et al; 2012). CPPA (Complex Polymeric Polyhydroxy Acid) is a group of organic acids that enhance various plant physiological functions such as nutrient absorption, shoot, and root growth. Figure 6 shows the visual comparison of seedling growth in different treatments

after 12 days of seed treatment application and seed sowing. These figures visualize the impact of different fertilizer products on seedling growth and development in Lentils.

7. Conclusion

This trial was conducted to assess the effectiveness of different fertilizer products on seed germination and seedling growth in lentils. The results revealed the application of Transit Zn and Momentum ZBM Trio significantly improved the shoot height. In addition, Transit Zn significantly increased the lentil root length. Therefore, it can be concluded that the application of Transit Zn as well as Momentum ZBM Trio as a seed treatment is highly beneficial in improving seed germination and seedling growth in lentils.

8. References

- Abares. (2007). Australian commodity statistics. Retrieved January 19, 2023, from <https://www.agriculture.gov.au/abares>
- Arshad, M., Adnan, M., Ahmed, S., Khan, A. K., Ali, I., Ali, M., ... & Khan, M. A. (2016). Integrated effect of phosphorus and zinc on wheat crop. *American-Eurasian Journal of Agriculture & Environmental Science*, 16(3), 455-459.
- Begum, M., Islam, M., Sarkar, M., Azad, M., Huda, A. and Kabir, A., 2016. Auxin signaling is closely associated with Zn-efficiency in rice (*Oryza sativa* L.). *Journal of Plant Interactions*, 11(1), pp.124-129.
- Graham, R. D., & Rengel, Z. (1993). Genotypic variation in zinc uptake and utilization by plants. *Zinc in Soils and Plants*, 107-118. doi:10.1007/978-94-011-0878-2_8
- Grains Research and Development Corporation. 2022. Our industry. [online] Available at: <<https://grdc.com.au/about/our-industry>> [Accessed 14 July 2022].
- Grant, C.A., and L.D. Bailey, 1989. Nitrogen, phosphorus, and zinc management effects on grain yield and cadmium concentration in two cultivars of durum wheat. *Canadian Journal of Plant Science*, 1: 63-70.
- Hajabbasi, M. and Schumacher, T., 1994. Phosphorus effects on root growth and development in two maize genotypes. *Plant and Soil*, 158(1), pp.39-46.
- Imran, M., Mahmood, A., Neumann, G., & Boelt, B. (2021). Zinc seed priming improves spinach germination at low temperatures. *Agriculture*, 11(3), 271. doi:10.3390/agriculture11030271
- Khan, A. S., Ahmad, B., Jaskani, M. J., Ahmad, R., & Malik, A. U. (2012). Foliar application of a mixture of amino acids and seaweed (*Ascophyllum nodosum*) extract improve growth and physicochemical properties of grapes. *Int. J. Agric. Biol*, 14(3), 383-388.
- Koger, C. H., Reddy, K. N., & Poston, D. H. (2004). Factors affecting seed germination, seedling emergence, and survival of texasweed (*Caperonia palustris*). *Weed science*, 52(6), 989-995.
- Montanha, G. S., Rodrigues, E. S., Marques, J. P., De Almeida, E., Colzato, M., & Pereira de Carvalho, H. W. (2020). Zinc nanocoated seeds: An alternative to boost soybean seed germination and seedling development. *SN Applied Sciences*, 2(5). doi:10.1007/s42452-020-2630-6
- OHKI, K., 1976. Effect of Zinc Nutrition on Photosynthesis and Carbonic Anhydrase Activity in Cotton. *Physiologia Plantarum*, 38(4), pp.300-304.
- Phuphong, P., Cakmak, I., Yazici, A., Rerkasem, B. and Prom-u-thai, C., 2020. Shoot and root growth of rice seedlings as affected by soil and foliar zinc applications. *Journal of Plant Nutrition*, 43(9), pp.1259-1267.
- Prom-u-thai, C., Rerkasem, B., Yazici, A., & Cakmak, I. (2012). Zinc priming promotes seed germination and seedling vigor of Rice. *Journal of Plant Nutrition and Soil Science*, 175(3), 482-488. doi:10.1002/jpln.201100332
- Talboys, P., Heppell, J., Roose, T., Healey, J., Jones, D. and Withers, P., 2015. Struvite: a slow-release fertiliser for sustainable phosphorus management?. *Plant and Soil*, 401(1-2), pp.109-123.

Xue, X., Du, S., Jiao, F., Xi, M., Wang, A., Xu, H., . . . Wang, M. (2021). The regulatory network behind maize seed germination: Effects of temperature, water, phytohormones, and nutrients. *The Crop Journal*, 9(4), 718-724. doi:10.1016/j.cj.2020.11.005

Sadras, V.O., Rosewarne, G.M. and Lake, L. (2021) "Australian lentil breeding between 1988 and 2019 has delivered greater yield gain under stress than under high-yield conditions," *Frontiers in Plant Science*, 12. Available at: <https://doi.org/10.3389/fpls.2021.674327>.

